

ROUTINE BLOODS for Peritoneal Dialysis Patients

JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC
UEC	UEC	UEC	UEC	UEC	UEC	UEC	UEC	UEC	UEC	UEC	UEC
Ca, PO4, Mg	Ca, PO4, Mg	Ca, PO4, Mg	Ca, PO4, Mg	Ca, PO4, Mg	Ca, PO4, Mg	Ca, PO4, Mg	Ca, PO4, Mg	Ca, PO4, Mg	Ca, PO4, Mg	Ca, PO4, Mg	Ca, PO4, Mg
FBC	FBC	FBC	FBC	FBC	FBC	FBC	FBC	FBC	FBC	FBC	FBC
LFT's	LFT's	LFT's	LFT's	LFT's	LFT's	LFT's	LFT's	LFT's	LFT's	LFT's	LFT's
Urate	Urate	Urate	Urate	Urate	Urate	Urate	Urate	Urate	Urate	Urate	Urate
			HBsAg (yearly if core positive) Anti-HBs						HepCAb HBsAg (yearly if core positive) Anti-HBs		
Tissue Typing*	Tissue Typing*	Tissue Typing*	Tissue Typing*	Tissue Typing*	Tissue Typing*	Tissue Typing*	Tissue Typing*	Tissue Typing*	Tissue Typing*	Tissue Typing*	Tissue Typing*
Troponin T*						Troponin T*					
Fe studies*			Fe studies*			Fe studies*			Fe studies*		
PTH			PTH			PTH			PTH		
HbA1c (only diabetic patients)			HbA1c (only diabetic patients)			HbA1c (only diabetic patients)			HbA1c (only diabetic patients)		
			Fasting lipids - Total Cholesterol, HDL, LDL & Triglycerides* only in high risk patients						Fasting lipids - Total Cholesterol, HDL, LDL & Triglycerides* only in high risk patients		
Tissue typing for patients on transplant list. Performed before 20th of each month and repeated 2 weeks after a blood transfusion.											
Fe studies performed 2 weeks after completing a course of IV Fe.											
HIV tesing is performed at commencement of dialysis.											
Fasting Lipids only for patients with a history of, or at risk of, CAD, CVD, PVD or diabetes.											
Troponin T : Twice per year											
Hepatitis B testing - refer to protocol. Core positive patients only require yearly antigen testing											
Blood Borne Viruses: Initial testing for HIV, Anti-HCV, HBsAg, Anti-HBs, Anti-HBc (informed consent required)											
Hepatitis B surface antibody (Anti-HBs) level 4-8 weeks post vaccination course											
Dialysis Adequacy (Kt/V and Ccl) - Annual tests											