

PRESERVE Study

Inclusion

- Planned/urgent angiography
- Diabetes with eGFR <60 ml/min
 - On insulin or oral medn only
- Anyone with eGFR <45 ml/min
 - Based on eGFR value taken in the last 1 month
 - No SCr in the last month? *SCr must be taken prior to enrolment THEN another for Study SCr must be taken*
- Ability to comply with D4 & D90 FUPs (endpoint assessments)
 - Patient to return to study hospital for D4 & D90 FUPs

Exclusion

- Emergent angiography <3hrs from identification to procedure
- Currently dialysis dependent
- eGFR <15 ml/min (Stage 5 CKD)
- Known unstable SCr values (defined as a change in SCr of $\geq 25\%$ over the 3 days prior to angiography)
- Decompensated heart failure requiring any of:
 - Inotropes, balloon pump, filtration
 - Not Nitrates
- Iodinated contrast in last 7 days
- Any NAC within last 48hrs
- Women Pregnant/breastfeeding
- Women (childbearing age) will not/cannot use adequate contraception
- Age <18
- Allergies/contraindications to study treatments or allergies to iodinated contrast (hives and/or trouble breathing)
- Unwilling/unable to provide consent



Contact

Jenny Brimley – Cardiac Clinical Trials Co-ordinator
Cardiology Department,
Level 1 Clinical Services Building,
Gray Street,

9113 3168
(or page 069)

Jennifer.brimley@sesiahs.health.nsw.gov.au